



## Fish Zoology

Sweazea, Simperova

Department of Zoological Research

Article Info

**Received: 16-12-2025 Revised: 11-1-2026 Accepted: 14-2-2026 Published: 19-3-2026**

### **Abstract**

Since its founding in 1984 as the continuation of Zoological Magazine (1888–1983) and *Annotationes Zoologicae Japonenses* (1897–1983), the Zoological Society of Japan, one of the country's oldest scientific societies, has published *Zoological Science*, a unique and prestigious international journal in zoology. The range of animal species that members of the Society and journal authors use in their research may be one of the most notable aspects of both organizations. Fish may have made contributions to nearly every field of study and publication, including behavioral biology, biochemistry, cell biology, developmental biology, diversity and evolution, ecology, endocrinology, genetics, immunology, morphology, neurobiology, phylogeny, reproductive biology, and taxonomy. Many fish species have been used as model animals in zoological science study due to the recent development of contemporary molecular genetic techniques in biology. The writers of this Special Issue, who range in age from junior to senior zoologists, have contributed a variety of works in zoological science using a range of fish species.

**Key words:** fish, neurobiology, physiology, morphology, genetics, and behavior biology

### **INTRODUCTION**

The range of animal species that members of the Zoological Society of Japan and its official journal, *Zoological Science*, use in their research may be one of the most notable aspects of both organizations. According to FishBase (<https://fishbase.mnhn.fr/search.php>), there are currently roughly 35,000 species of fish, or about half of all vertebrates. Because fish species are so diverse, they have a wide range of morphological, behavioral, and developmental characteristics, some of which are particularly useful for research with particular goals. Fish may have contributed to nearly every discipline in presentations and publications in *Zoological Science*, and several investigations have been carried out by utilizing these qualities of fish. Many fish species have been used as model animals in zoological science study due to the recent development of contemporary molecular genetic methods in biology. As of February 2023, a search on the ZooDiversity website (<https://zdw.zoology.or.jp/>) revealed that over 1000 publications published in *Zoological Science* have employed fish. The writers of the current Special Issue, who range in age from junior to senior zoologists, have contributed a variety of fish-based zoological research works. This issue includes GENETICS (1), PHYSIOLOGY (1), MORPHOLOGY (1), NEUROBIOLOGY (4), and BEHAVIORAL BIOLOGY (2). It is composed of a balanced mix of five review papers and four original studies. Eight of these nine publications concentrate on teleost fishes (medaka, goby, goldfish, electric fish, guppy, cichlid fish, and so forth), while one article focuses on research employing cartilaginous fish.

### **GENETICS**

The authors of the paper by Inoue and Takeda (2023) examine Teratorn and its associated elements, a unique class of herpesviruses that are common in teleost genomes. Inoue et al. (2017) discovered a new class of herpesviruses called Teratorn and its associated components in the genomes of several teleost fish species. A

piggyBac-like DNA transposon is fused to at least part of the Teratorn-like herpesviruses, indicating that they have taken over the transposon-like intragenomic lifestyle by abusing the trans-poson system. Inoue and Takeda (2023) talked about the life cycle, the generality of transposon-virus fusion, and the sequencing features of Teratorn-like herpesviruses as well as their evolutionary ties with other herpesviruses. The authors contend that Teratorn-like herpesviruses offer hard proof that non-retroviral components can develop into intragenomic parasites with the ability to replicate by obtaining transposition machinery from other sources. The first example is the herpesviruses that resemble Teratorn of herpesvirus adaptation to the intragenomic life cycle, and the research may contribute to the understanding of the evolutionary mechanisms of “jumping genes”, which may play important roles in species diversifications and evolution.

### **PHYSIOLOGY**

In the article by Horie et al. (2023), the authors focused on adaptation to a high salinity marine environment exhibited by cartilaginous fishes, which have evolved a highly elaborate urea-based osmoregulation (ureosmotic) strategy called “four-loop nephron” in the kidney. The characteristic structure of the “four-loop nephron” has been proposed to play important roles in performing dedicated roles in each segment and contribute to its overall function, such as urea reabsorption from the glomerular filtrate to maintain a high concentration of urea in the body (Hyodo et al., 2014).

This mechanism is considered to be important for reabsorption of urea from glomerular filtrate to maintain a high concentration of urea in the body. Horie et al. (2023) performed laser microdissection (LMD) of the kidney of cloudy catshark, *Scyliorhinus torazame*, and RNA-sequencing (RNA-seq) analysis to obtain segment-dependent gene expression profiles. They further examined expressions of the Na<sup>+</sup>-coupled cotransporters abundantly expressed in the second loop samples. Although the proximal II (PII) segment of the second loop is known for the elimination of excess solutes, the authors' results suggest that the PII segment is also crucial for reabsorption of valuable solutes. Since the regulatory mechanisms of renal function currently remain poorly understood in euryhaline elasmobranchs, the segment-dependent gene expression profiling reported here is expected to be a powerful method for unraveling the renal mechanisms and regulation in euryhaline elasmobranchs.

### **MORPHOLOGY**

In the article by Hagio and Yamamoto (2023), the authors review the ascending visual pathways from the retina to the telencephalon in teleosts, focusing on forebrain visual centers, associated neural circuitries, and evolution. Most vertebrates, including fish, recognize food and predators for survival, as well as individuals of the opposite sex for reproduction, by using various sensory systems such as the visual sensory system. From the authors' recent studies (e.g., Hagio et al., 2021), they propose that, in the common ancestor of teleosts, there were two visual pathways, the geniculate-like or thalamo-fugal pathway and the extrageniculate-like or tecto-thalamo-fugal pathway, and the geniculate-like system was lost in acanthopterygian fishes, which emerged relatively recently during evolution. Their in-depth analyses on the connections of visual centers also revealed that there are connections shared with those of mammals, and retinotopic organization of the ascending connections is maintained at least to the level of the diencephalon in the acanthopterygian teleosts, such as the yellowfin goby.

The authors also propose neural circuitries involving telencephalic visual centers and their view of the evolution of highly specialized compartments of the telencephalic visual receptive zone (lateral part of dorsal telencephalon: Dl) found in gobiid fish.

## NEUROBIOLOGY

In the article by Kawasaki (2023), the author reviews the time-comparator neural circuits of gymnotiform electric fishes. A major goal of behavioral neuroscience is to explain functions of sensory systems in terms of activity, connectivity, and computational features of peripheral and central neurons that are involved in the control of behavior. In sensory systems, information is typically encoded in the number of action potentials per unit time (rate coding) or in the firing timing or time patterns of action potentials (time coding). While identifying which coding scheme is used at a particular level of neuronal circuits is generally difficult due to the complex relations between sensory input signals and action potentials responding to them, the electrosensory system of gymnotiform electric fishes offers an exceptional substrate where the time-coding neural pathway is distinguished from rate-coding pathway with exceptional clarity. Kawasaki (2009) previously published a comprehensive review of electrosensory systems including the rate-coding pathways in *Zoological Science*. This time (Kawasaki, 2023), the author focused on comparative anatomy and physiology of the timing pathway in a gymnotiform pulse-type fish, *Brachyhypopomus*, and a gymnotiform wave-type fish, *Apteronotus*, in reference to the previous findings on other species. The author also discusses the timing system of mormyriiform electric fishes, which had evolved electrogenesis and electrosense independently from gymnotiform electric fishes. He concludes that comparative studies of the timing system of gymnotiform fishes have revealed neural circuits that are central for the detection of microsecond time differences between sensory signals with behavioral significance.

In the article by Oka (2023), the author reviews the neural control of sexual behavior in fish. Many vertebrate species show breeding periods and exhibit series of characteristic species-specific sexual behaviors only during the breeding period. Here, secretion of gonadal sex hormones from the mature gonads has been considered to facilitate sexual behaviors. Thus, the sexual behavior has long been considered to be regulated by neural and hormonal mechanisms. The author discusses recent progress in the study of neural control mechanisms of sexual behavior with a focus on studies using fish, which have often been the favorite animals used by many researchers who study instinctive animal behaviors. First, the control mechanisms of sexual behaviors by sex steroids are discussed in relation to the anatomical studies of sex steroid-concentrating neurons in various vertebrate brains, which are abundantly distributed in evolutionarily conserved areas such as preoptic area (POA) and anterior hypothalamus. The next focus is another brain area called the ventral telencephalic area, which has also been suggested to contain sex steroid-concentrating neurons and has been implicated in the control of sexual behaviors, especially in teleosts (Satou et al., 1984). The control of sex-specific behaviors and sexual preference influenced by estrogenic signals or by olfactory/pheromonal signals is also discussed. Finally, research on the modulatory control of motivation for sexual behaviors by a group of peptidergic neurons called terminal nerve (TN)-gonadotropin releasing hormone (GnRH) neurons (Umatani et al., 2022), which are known to be especially developed in fishes among various vertebrate species, is discussed.

In the article by Ikenaga et al. (2023), the authors performed histological and molecular characterization of the inferior olivary nucleus and climbing fibers, one of the main afferent input pathways to cerebellum, the goldfish.

The cerebellum in all vertebrates receives two main inputs: mossy fibers originating from a variety of precerebellar nuclei within the spinal cord and brainstem, and 2)

climbing fibers from the contralateral inferior olivary nucleus in the ventral medulla. The climbing fibers in mammals are known to entwine and terminate onto both major and peripheral branches of dendrites of the Purkinje cells.

However, the inferior olivary nucleus in teleost fishes has not been characterized well. Furthermore, previous neuroanatomical studies in teleosts failed to show whether climbing fibers only contact Purkinje cells or whether they also terminate on distal dendrites of eurydendroid cells, another type of cells in the teleost cerebellum. Ikenaga et al. (2023) first identified inferior olivary neurons by retrograde neural tracer application to the cerebellar hemisphere and found that most of them are calretinin-immunoreactive. Double labeling immunofluorescence with anti-calretinin and anti-zebrin II (marker of Purkinje cells; Ikenaga et al., 2005) antisera revealed that the calretinin-ir climbing fibers made synaptic-like contacts on the major dendrites of the zebrin II-ir Purkinje cells. These results support the idea that Purkinje cells, but not eurydendroid cells, receive strong inputs via the climbing fibers in teleosts, similar to the mammalian situation.

In the article by Tanaka et al. (2023), the authors used the whole cell patch clamp method to analyze firing activities of the TN-GnRH neurons in medaka during application of acetylcholine (ACh), which is one of the essential neuromodulators in the brain. Vertebrates generally possess hypophysiotropic and non-hypophysiotropic GnRH neurons. The TN-GnRH neurons are known to belong to the non-hypophysiotropic neurons and have been suggested to modulate sexual behaviors. These neurons show spontaneous pacemaker firing activity and release neuropeptides GnRH and neuropeptide FF to modulate motivation for male sexual behavior (Umatani et al., 2022). Since the spontaneous firing activities of peptidergic neurons, including GnRH neurons, are believed to play important roles in the release of neuropeptides, understanding the regulatory mechanisms of these spontaneous firing activities is important. The authors' results demonstrated that ACh induces hyperpolarization and inhibits their pacemaker firing. Electrophysiological analysis using an antagonist for acetylcholine receptors and *in situ* hybridization analysis showed that firing of TN-GnRH neurons is inhibited via M2-type muscarinic acetylcholine receptor. Taken together with literature from several other fish species including teleosts and elasmobranchs, the authors conclude that ACh may generally play an inhibitory role in modulating spontaneous activities of TN-GnRH neurons and thereby sexual behaviors in fish.

## **BEHAVIORAL BIOLOGY**

In the article by Takeuchi (2023), the author reviews the developmental process of a pronounced laterality in the scale-eating cichlid fish *Perissodus microlepis* in Lake Tanganyika. Using high-speed video recordings, Takeuchi et al. (2012) previously showed that predation behavior of *Perissodus microlepis* consists of five subcomponents. Further, in a preferred-side attack, in which the mouth-opening direction coincides with the attack direction, the fish can bend its body more rapidly and strongly at the time of predation than in a non-preferred-side attack, increasing the chances of a successful predation. These experiments provided direct evidence that the morphological differences in mouth structure and kinetic differences in bending movements allow the scale-eating fish to engage in lateral predatory behavior. The authors' recent studies have also shown that behavioral laterality in this fish depends on both genetic factors and past experience.

The attack-side preference of scale eaters is an acquired trait developed in the early developmental stage. Juvenile fish empirically learn which side of the prey is more effective for tearing scales and gradually select the dominant side for attacking. However, the superior kinetics of body flexion during the dominant side attack has innate characteristics that are not altered by experience. Additionally, left-right differences in scale-

eater mandibles also develop during ontogeny.

The author concludes by suggesting that further progress toward understanding the comprehensive mechanisms of laterality should address the following persistent barriers: (1) the effects of phylogenetic constraints and ecological factors on the level of laterality, and (2) the neuronal and molecular mechanisms that produce left–right behavioral differences. In the article by Sogawa et al. (2023), the authors investigated the visual signal for individual-recognition in males of sexually dichromatic guppy (*Poecilia reticulata*, Cyprinodontiformes). Individual recognition is a necessary cognitive ability for the maintenance of stable social relationships. Recent studies have shown that, like primates, some fish species can distinguish familiar fish from unfamiliar strangers via face-recognition. However, the taxa of the studied fish species are restricted to Perciformes, and the visual signal used for the recognition of fish remains unclear. Kohda et al. (2015) reported on face-recognition by exploiting small individual differences in the shape and size of color patches near the eyes on the operculum in the territorial cichlid *Neolamprologus pulcher*. Using male guppies, Sogawa et al. (2023) examined the hypothesis that fish distinguish between familiar individuals and unknown strangers by their faces rather than by body coloration.

Here, the authors made an important finding that male faces contain clear individual-variation in white/metallic colored patches on the operculum visible to the human eyes. From the fact that the present photo-model eliminated UV-light information (note that guppies are known to be capable of UV vision), they suggest that these patches might be an important visual stimulus for face-recognition in male guppies, like some cichlids. From the comparison among males of different guppy variants, including wild-type phenotype, the authors suggest that the face color-patch is stable regardless of variation in body color, with a different genetic mechanism potentially underlying face and body colors. As described above, the present Special Issue covers many disciplines of research fields in zoological science by

taking advantage of various fish species. We hope that the readers enjoy reading the entire issue and take this opportunity to discuss the future of the research in zoological science as a whole.

## REFERENCES

- Hagio H, Kawaguchi M, Abe H, Yamamoto N (2021) Yellowfin goby *Acanthogobius flavimanus*: Afferent and efferent connections of the prethalamic nucleus. *J Comp Neurol* 529: 87–110
- Ascending visual routes to the telencephalon in teleosts with particular attention on forebrain visual centers, related neural circuitries, and evolution (Hagio H, Yamamoto N, 2023). *Zool Sci* 40: 105–118
- Horie T, Takagi W, Aburatani N, Yamazaki M, Inokuchi M, Tachizawa M, et al. (2023) Laser microdissection for segment-level functional characterisation of the cartilaginous fish nephron using segment-dependent gene expression profiling. *Zool Sci* 40: 91–104
- Hyodo S, Kakumura K, Takagi W, Hasegawa K, Yamaguchi Y (2014) Kidney morphology and function in cartilaginous fishes: a focus on urea reabsorption. *Am J Physiol Regul Integr Comp Physiol* 307: R1381–R1395
- Ikenaga T, Yoshida M, and Uematsu K (2005) Efferent neurons of the goldfish corpus cerebelli: morphology and immunohistochemistry. *J Comp Neurol* 487: 300–311
- Ikenaga T, Morita S, Finger TE (2023) Histological

**Frontiers in Zoological Research**  
**Volume 2 Issue 1 2026**

and molecular analysis of the climbing fibers and inferior olivary nucleus in the goldfish, *Carassius auratus*. *Zool Sci* 40: 83–90 Inoue Y, Takeda H (2023) Teratorn and its associated elements—a novel category of herpesviruses common in teleost genomes—*Zool Sci* 40: 141–150

The Teratorn mobile element in medaka fish was produced by the full fusion of a transposon and herpesvirus (Inoue Y, Saga T, Aikawa T, Kumagai M, Shimada A, Kawaguchi Y, et al., 2017). Kawasaki M. (2009) Evolution of time-coding systems in weakly electrified fishes, *Nat Commun* 8: 551. *Zool Sci* 26: 587–599 Kawasaki M. (2023) Gymnotiform electric fish time-comparator brain circuits. *Zool Sci* 40: 119–127 Kohda M, Jordan LA, Hotta T, Kosaka N, Karino K, Tanaka H, et al. (2015) Facial recognition in a cichlid fish that lives in a community. Neural regulation of fish sexual behavior, *PLoS One* 10: e0142552 Oka Y (2023). *Zool Sci* 40: 128–140

Satou M, Oka Y, Kusunoki M, Matsushima T, Kato M, Fujita I, et al. (1984) Results of electrical brain stimulation tests show that telencephalic and preoptic areas integrate sexual behavior in hime salmon (landlocked red salmon, *Oncorhynchus nerka*). 33: 441–447 *Physiol Behavior* Male guppies identify familiar conspecific males by their faces (Sogawa S, Fukushima R, Sowersby W, Awata S, Kawasaka K, Kohda M, 2023). *Zool Sci* 40: 168–174

Takeuchi Y (2023) described how the scale-eating cichlid fish *Perissodus microlepis* in Lake Tanganyika developed a noticeable laterality. *Zool Sci* 40: 160–167

Lateralized kinematics of predation behavior in a Lake Tanganyika scale-eating cichlid fish (Takeuchi Y, Hori M, Oda Y, 2012). *PLoS One* 7: e29272